

Derivatives of Exponential, Logarithmic Functions

Given:

if $y = e^x$, then $\frac{dy}{dx} = \underline{\hspace{2cm}}$ Answer: e^x

if $y = \ln x$ then find $\frac{dy}{dx} = \underline{\hspace{2cm}}$ Answer: $\frac{1}{x}$

ex1 If $y = xe^x - e^x - 2x^2$, use algebra to determine any relative maximum or minimum values. Is there an absolute minimum or maximum value?

(Answer: local max (0,-1), local min ($\ln 4$, $4\ln 4 - 2 - 2(\ln 4)^2$) which is about -2.30)

Ex2 Find the equation of the tangent to

$$f(x) = \frac{x}{e} \ln\left(\frac{e}{x}\right) \quad (\text{Answer: } y = \frac{-2}{e}x + 2)$$

when

$$x = e$$

Ex 3 Suppose $h = e^{\ln t} - \ln(t^2 + 1)$. (t in seconds, h in metres). When is the velocity equal to zero? Is this a maximum or minimum height? Find the acceleration. Is there a minimum or maximum velocity?

Answer: a) at $t=1$ b) neither, as the velocity has a double root at $t=1$

c) $a = \frac{2(t^2 - 1)}{(t^2 + 1)^2}$ d) yes, at $t=1$ the acceleration goes from - to +, so min v at $t=1$.

Ex 4 Find $\frac{dy}{dx}$ at (1,0) if $\frac{(\ln x)^2}{e^y} - \ln(x+y) = e^{\sin(2x-2)\cos^2 y} - 1$

(Answer: $\frac{dy}{dx} = -3$)

Ex5 Find the derivative of $y = 5^x$. Then try to find a general result.

Answer: a) $\frac{dy}{dx} = (\ln 5) \cdot 5^x$ and in general, $\frac{dy}{dx} = (\ln a) \cdot a^x$, where $y = a^x$

Ex6 Find the derivative of $y = \log_3 x$. Then try to find a general result.

Answer: $\frac{dy}{dx} = \frac{1}{\ln 5} \left(\frac{1}{x} \right)$, and in general, $\frac{dy}{dx} = \frac{1}{\ln a} \left(\frac{1}{x} \right)$, where $y = \log_a x$

Ex 7 How would you find the derivative of $y = x^x$? What is the minimum value of this function?

(Answer: Take ln of both sides first and then take the derivative)

min value is $\left(\frac{1}{e} \right)^{\frac{1}{e}} \doteq 0.692$

Ex 8 Find the smallest positive value of the constant k such that the graphs of $y = e^x$ and $y = k \sin x$ are tangent to one another. Find also the point of tangency.

(Answer: $k = \sqrt{2}e^{\frac{\pi}{4}} \doteq 3.10$ and point is $\left(\frac{\pi}{4}, e^{\frac{\pi}{4}}\right)$)

Exponential /Logarithmic Functions- Limits

Exponential limits may still include l'Hopital's rule, order of size, using a substitution and first principles of derivatives as methods of solution, but there are two new types of limits we will discuss involving **definition of e^x and logarithmic method.**

Ex 1 Evaluate the following limits:

$$\text{a) } \lim_{x \rightarrow \infty} \frac{2^x + x}{2^{x+1} + x^2}$$

$$\text{b) } \lim_{x \rightarrow \pi} \frac{\ln(\sin \frac{x}{2})}{\cos x + 1}$$

$$\text{c) } \lim_{x \rightarrow \infty} x \ln\left(\frac{1}{x} + 1\right)$$

$$\text{d) } \lim_{x \rightarrow \infty} \left(1 + \frac{1}{2x}\right)^x$$

$$\text{e) } \lim_{x \rightarrow \infty} x\left(1 - \frac{1}{x}\right)^x$$

$$\text{f) } \lim_{x \rightarrow \infty} 4xe^{1/x} - 4x$$

Answers: a) 0 b) $-\frac{1}{4}$ c) 1 d) \sqrt{e} e) no limit f) 4

Day 9- Problems- Logarithms, Exponential Functions

1. Solve for x without a calculator. Leave answers in exact form:

a) $6 + e^x = 16e^{-x}$ b) $\ln x + \log_{e^2} x = -3$

2. Find the equation of the tangent to $y = 3^{2x-2} - 3^{x-1} - (\ln 3)x + \ln 3 + 2x$ at $x=1$.
Leave answers in exact form.

3. Find the value of the constant k such that the graphs of

$y = \ln x$ and $y = -\frac{1}{3}x^3 + k$ will intersect at right angles.

4. The graphs of $f(x) = x^2$ and $g(x) = k \ln x$, where k is a positive constant are tangent to one another. Find the point of tangency and the exact value of k .

5. If $e^{xy} = 2$, then at the point $(1, \ln 2)$, $\frac{dy}{dx} =$

a) $-\ln 2$ b) $2\ln 2$ c) $\ln 2$ d) $-2e$ e) $-4\ln 2$

6. If $f(x) = e^{-x} + 2$ for $x < 0$ and $f(x) = ax + b$ for $x \geq 0$, and if $f(x)$ is differentiable at $x=0$, then $a+b=$

a) 0 b) 1 c) 2 d) 3 e) 4

7. Evaluate: a) $\lim_{x \rightarrow \infty} e^x \ln(1-x)$ b) $\lim_{x \rightarrow 0} (2 \cos x - 1)^{\frac{1}{x}}$

8.

Find any value of k , where k is a constant, such that the graphs of

$y = k^x$ and $y = \log_k x$ will be tangent to one another. Find the point of tangency.

(Solution follows on next page)

Answers: 1. a) $\ln 2$ b) $\frac{1}{e^2}$ 2. $y = 2x$ 3. $k = \frac{1}{3}$ 4. point (\sqrt{e}, e) and $k = 2e$

5. A 6. C (a=-1, b=3) 7. a) 0 b) 1

Full Solution to #8

$$y = \log_k x \quad \text{eq1} \quad \Leftrightarrow y = \frac{\ln x}{\ln k}$$

$$y = k^x \quad \text{eq2}$$

$$\text{let } k^x = \frac{\ln x}{\ln k} \quad \text{eq3}$$

also, slopes are equal,

$$\therefore \ln k(k^x) = \frac{1}{x \ln k} \Leftrightarrow k^x = \frac{1}{x(\ln k)^2} \quad \text{eq4}$$

Equating,

$$\frac{1}{x(\ln k)^2} = \frac{\ln x}{\ln k}$$

$$\ln k = \frac{1}{x \ln x} \Rightarrow k = e^{1/(x \ln x)} = e^{(x \ln x)^{-1}}$$

But in order to maximise k , let $\frac{dk}{dx} = 0$

$$0 = e^{1/x \ln x} (-1)(x \ln x)^{-2} [\ln x + 1]$$

$$\Rightarrow \ln x + 1 = 0 \Rightarrow x = \frac{1}{e}$$

\therefore the point of tangency is $\left(\frac{1}{e}, e^{1/(\ln x)} \right)$

